




REVISTA DE
GASTROENTEROLOGÍA
DE MÉXICO

www.elsevier.es/rgmx



CLINICAL IMAGE IN GASTROENTEROLOGY

An uncommon gallbladder location associated with polycystic liver[☆]



Una localización poco frecuente de la vesícula biliar asociada con el hígado poliquístico

V.J. Alegre Salles*, C.E. Azevedo Ferreti, M.M. Natrielli de Almeida

Servicio de Cirugía General y Digestiva, Hospital Regional Vale del Paraíba, Taubaté, San Pablo, Brazil

The gallbladder is located between segments IV and V of the liver in a bare area, not covered by the visceral peritoneum.



Figure 1 Abdominal computed tomography scan showing voluminous hepatic cysts.

An 81-year-old woman presented with chronic abdominal pain in the epigastrium and right hypochondrium, associated with weight loss and an abdominal tumor. A CT scan revealed a polycystic liver with a massive hepatic cyst in the right lobe (**fig. 1**).

Surgical exploration identified the gallbladder surrounded by the thick serosal layer of a hepatic cyst (**fig. 2**).

The atypical location of the gallbladder alerted the surgical team to the possibility of a biliary tract abnormality, which was ruled out after performing an intraoperative cholangiography. Polycystic liver is estimated to occur in approximately 5% of the population. Exaggerated growth of the cysts can cause abdominal pain and distension, dyspepsia, early satiety, dyspnea, and a visible or palpable




Figure 2 Subcapsular position of the gallbladder.

[☆] Please cite this article as: Alegre Salles VJ, Azevedo Ferreti CE, Natrielli de Almeida MM. Una localización poco frecuente de la vesícula biliar asociada con el hígado poliquístico. *Revista de Gastroenterología de México*. 2015;80:109–110.

* Corresponding author at: University of Taubaté, Medicine, Avenue Tiradentes, 280 Taubaté, São Paulo 12030180 Brazil.
Phone: +550121536813888; Home: +550121536316061;
fax: +55011536316061.

E-mail address: geriaclinsecretaria@gmail.com
(V.J. Alegre Salles).

abdominal tumor. The cysts do not usually reach the biliary or vascular structures, but if that occurs, it can compress the biliary tract and cause jaundice in up to 9% of the patients.

Ethical responsibilities

Protection of persons and animals. The authors declare that no experiments were performed on humans or animals for this study.

Data confidentiality. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Financial disclosure

No financial support was received in relation to this study/article.

Conflict of interest

The authors declare that there is no conflict of interest.